

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/355504543>

Fruit and vegetable intake and mental health among family caregivers of people with dementia in Uganda

Article in *Mental Health and Prevention* · October 2021

DOI: 10.1016/j.mhp.2021.200223

CITATIONS

0

READS

41

7 authors, including:



Herbert Ainamani

Kabale University

15 PUBLICATIONS 46 CITATIONS

[SEE PROFILE](#)



Godfrey Zari Rukundo

Mbarara University of Science & Technology (MUST)

134 PUBLICATIONS 709 CITATIONS

[SEE PROFILE](#)



Sam Tumwesigire

Kabale University

7 PUBLICATIONS 104 CITATIONS

[SEE PROFILE](#)



Alexander C Tsai

Massachusetts General Hospital

372 PUBLICATIONS 11,927 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



CHAKA Project [View project](#)



Article Prevalence Of Violence-Related Traumatic Events And Symptoms Of Post-Traumatic Stress Disorder Among The Congolese Refugees In Uganda [View project](#)



Fruit and vegetable intake and mental health among family caregivers of people with dementia in Uganda

Herbert E. Ainamani^{a,*}, Wilson M. Bamwerinde^b, Godfrey Z. Rukundo^c, Sam Tumwesigire^d, Valence Mfitumukiza^e, Everd M. Bikaitwoha^d, Alexander C. Tsai^f

^a Department of Mental Health, Kabale University School of Medicine, Kabale, Uganda, P. O. Box 317

^b Department of Environment and Natural Resources Kabale University, Kabale, Uganda

^c Department of Psychiatry, Mbarara University of Science and Technology, Mbarara, Uganda

^d Department of Pediatrics, Kabale University School of Medicine, Kabale, Uganda

^e Department of Nursing, Kabale University School of Medicine, Kabale, Uganda

^f Center for Global Health and Mongan Institute, Massachusetts General Hospital, Boston, United States

ARTICLE INFO

Keywords:

Anxiety
Dementia
Depression
Diet
Mental health
Stress
Sub-Saharan Africa
Uganda

ABSTRACT

Consumption of fruits and vegetables is correlated with improved mental wellbeing. Although this growing body of research has been recognized by researchers and clinicians in high-income countries, fewer studies examining this relationship have been conducted in low- and middle-income settings. In this study, we sought to estimate the association between fruit and vegetable intake and symptoms of depression and anxiety. We conducted a cross-sectional study among 242 family caregivers of people with dementia in southwestern Uganda. Fruit and vegetable intake in the past week was measured with a food frequency questionnaire. Depression and anxiety were assessed using the depression and anxiety subscales of the 42-item Depression, Anxiety and Stress Scales. Multivariable regression models were used to estimate the associations between fruits and vegetable consumption and depression and anxiety, adjusting for caregiving burden and other potential confounders. Depression symptom severity was negatively associated with consumption of jackfruits ($b = -4.68$; 95% confidence interval [CI], -8.96 to -0.39), green leafy vegetables ($b = -14.1$; 95% CI, -18.0 to -10.1), root vegetables ($b = -14.0$; 95% CI, -19.5 to -8.63), and other vegetables ($b = -14.8$; 95% CI, -19.3 to -10.3), and frequent consumption of vegetables ($b = -1.91$; 95% CI, -3.77 to -0.04). Anxiety symptom severity was negatively associated with consumption of green leafy vegetables ($b = -12.2$; 95% CI, -16.0 to -8.46), root vegetables ($b = -12.6$; 95% CI, -17.5 to -7.58), and other vegetables ($b = -12.7$; 95% CI, -17.0 to -8.40), and frequent consumption of vegetables ($b = -2.07$; 95% CI, -3.84 to -0.29). Our results suggest that fruit and vegetable consumption is associated with reduced depression and anxiety symptoms.

1. Background

Depression and anxiety are the most commonly studied mental health problems among the caregivers of older-age people with dementia (Alfakhri et al., 2018; Clyburn, Stones, Hadjistavropoulos, & Tuokko, 2000; Omranifard, Haghighizadeh, & Akouchekian, 2018), people with stroke (Alfakhri et al., 2018; Atteih et al., 2015; Balhara, Verma, Sharma, & Mathur, 2012; Guo & Liu, 2015) and those with mental ill health (Alqahtani et al., 2018; Minichil, Getinet, Derajew, & Seid, 2019). Studies on dementia report that, in addition to experiencing high levels of caregiving burden, caregivers of older-age people with

Alzheimer's disease and other related dementias (ADRD) tend to have symptoms of depression and anxiety (Clyburn et al., 2000; Dotchin et al., 2014). A meta-analysis that included 17 studies on the prevalence of mental health disorders among caregivers of patients with Alzheimer's disease found that more than one-third had clinically significant symptoms of depression and anxiety (Sallim, Sayampanathan, Cuttilan, & Ho, 2015). In fact, the authors of the above meta-analysis observed that when compared with caregivers of other diseases, the prevalence of mental health problems among caregivers of people with dementia was higher than caregivers of other conditions.

Recent studies in nutrition and mental health, mostly from high

* Corresponding author.

E-mail addresses: hainamani@kab.ac.ug (H.E. Ainamani), wbamwerinde@kab.ac.ug (W.M. Bamwerinde), grukundo@must.ac.ug (G.Z. Rukundo), s.tumwesigire@kab.ac.ug (S. Tumwesigire), smfitumukiza@kab.ac.ug (V. Mfitumukiza), ebmaniple@kab.ac.ug (E.M. Bikaitwoha), actsai@mg.harvard.edu (A.C. Tsai).

<https://doi.org/10.1016/j.mhp.2021.200223>

Received 30 January 2021; Received in revised form 11 October 2021;

Available online 21 October 2021

2212-6570/© 2021 The Authors. Published by Elsevier GmbH. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

income countries, suggest that consumption of some foods may be associated with improvements in mood (Gilbody, Lightfoot, & Sheldon, 2007; Li, Li, Song, & Zhang, 2017; McMartin, Jacka, & Colman, 2013; Péneau et al., 2011; Van der Does, 2001). For example, higher fruit and vegetable intake has been found to be associated with lower symptoms of depression and anxiety (Gibson-Smith et al., 2020; Liu, Yan, Li, & Zhang, 2016). In a cross-sectional study among 400 women with mental health problems attending health care centers, it was found out that participants in the lower quartile of fruit and vegetable intake were more likely to develop depression than those taking higher doses (Baharzadeh et al., 2018). Consistent with the above literature, a large meta-analysis of eighteen studies on fruit and vegetable intake observed a negative correlation between the consumption of fruits/vegetables and depression (Liu et al., 2016).

Although previous research studies in high-income countries have linked increased fruit and vegetable intake to lower levels of depression (McMartin et al., 2013; Tsai, Chang, & Chi, 2012) and anxiety (McMartin et al., 2013; Smith & Rogers, 2014), few studies about this phenomenon have been conducted in sub-Saharan Africa. One national survey of 3962 adult participants showed that the consumption of both fruits and vegetables in Uganda is low (Kabwama, Bahendeka, Wesonga, Mutungi, & Guwatudde, 2019). Yet the mental health benefits of fruit and vegetable consumption are well documented (Liu et al., 2016).

Given the significant role family caregivers play in the care of older-age persons with ADRD in Africa (Ainamani et al., 2020b; Kakongi et al., 2020; Kamoga et al., 2019; Mubangizi, Maling, Obua, & Tsai, 2020) and their high levels of symptoms depression and anxiety (Ainamani et al., 2020a; Ainamani et al., 2021), it is important to provide evidence on potentially modifiable factors that can reduce the mental health burden of depression and anxiety in this particular group of caregivers. Our study aimed to estimate the association between fruit and vegetable intake and symptoms of depression and anxiety in a sample of family caregivers of people with ADRD in a rural area of southwestern Uganda. We hypothesized that the consumption of different types of fruits and vegetables would be negatively correlated with symptoms of depression and anxiety among the caregivers of people living with dementia in rural, southwestern Uganda.

2. Methods

2.1. Population and design

We conducted a cross-sectional study to assess fruit and vegetable consumption and symptoms of depression and anxiety among 242 adult family caregivers of patients with ADRD in two districts of Kigezi, a rural region of southwestern Uganda. We collected data during July-August 2020. To be included in the study, participants had to be 18 years of age or older, identify as the family caregiver of someone living with ADRD, live in the same house or compound with the patient, and have provided care for more than 6 months. (In the remainder of this manuscript, for ease of exposition we refer to the caregivers as the “study participants” and to the family members with ADRD as their “patients.”) We sought guidance from the health workers of Reach One Touch One Ministries (ROTOM) (Rukiga District) and Heal Medical Centre (Rubanda District) to locate family caregivers of people with ADRD. We excluded caregivers who could not communicate to our study staff reliable information about their patients (e.g., due to deafness/mutism or acute intoxication). All interviews were conducted in Runyankore-Rukiga (the local language).

2.2. Ethical considerations

Approval to conduct this research was obtained from the Mbarara University of Science and Technology Research Ethics Committee, and clearance was obtained from the Uganda National Council for Science and Technology. Before the interview, information on the content,

procedures, risks, the right to withdraw, and confidentiality were explained to prospective participants. Both oral and written informed consent were provided by each research participant at the beginning of every interview. Participants who did not know how to read and write were asked to provide a thumbprint for consent.

2.3. Primary exposures and outcomes of interest

Fruit and vegetable consumption was assessed using a dietary survey questionnaire (Brookie, Best, & Conner, 2018) that assesses the weekly consumption of different fruits and vegetables. For each fruit and vegetable type, study participants were asked the number of days per week of consumption (ranging from 0-7) and the number of servings consumed per day (ranging from 1-7 servings). Frequency of weekly milk consumption was assessed similarly. Specification of these variables was based on previous work (Baharzadeh et al., 2018; Choda et al., 2020; Crichton, Murphy, & Bryan, 2010).

The 42-item Depression, Anxiety and Stress Scales (DASS-42) was used to assess caregivers' symptoms of depression and anxiety (Lovibond & Lovibond, 1995). This scale consists of depression, anxiety, and stress subscales, each containing 14 items. The items in each subscale are scored on a 4-point Likert-type scale (“never”, “sometimes”, “often”, “almost always”), yielding a total score ranging from 0 to 42 for each subscale, with higher scores indicating more severe symptoms of depression, anxiety, and stress. The DASS-42 has previously shown adequate internal consistency, with evidence of construct validity suggested by robust correlations with associated constructs (Ainamani et al., 2020a; Crawford & Henry, 2003; Gloster et al., 2008). In our sample, the Cronbach's α was 0.89.

2.4. Covariates

In addition to measuring the main explanatory variables, caregiving burden was assessed using the Zarit Burden Interview (ZBI) (Zarit, Reever, & Bach-Peterson, 1980). The ZBI is a 22-item scale that assesses various aspects of caregiving burden. The items are scored on a 5-point Likert scale that ranges from “never” to “nearly always,” with a total sum score that ranges between 0 and 88. This scale is commonly used in research studies of caregiving burden and its association with mental health problems (Ainamani et al., 2020a; Springate & Tremont, 2014). Patient functional status was measured using the Bristol Activities of Daily Living Scale (BADLS) (Bucks, Ashworth, Wilcock, & Siegfried, 1996). BADLS total scores range from 0 to 60, with higher scores indicating more severe functional impairment (Ainamani et al., 2020a; Jefferson et al., 2008). We also measured additional covariates: age and sex of both caregivers and their patients, the nature of the patients' relationships with their caregivers, total years of caregiving, and caregivers' educational attainment in years. Inclusion of these covariates was based on previous work linking them to mental health problems (Ainamani et al., 2020a; Kwon, Park, Kim, Choi, & Jang, 2019).

2.5. Data analysis

Using Stata software (version 16, Stata Corp., College Station, Tex.), we computed standard statistics to summarize characteristics of the sample. The association between fruit and vegetable consumption and mental health outcomes (depression and anxiety subscale scores specified as continuous variables) was estimated using multivariable linear regression models. To assess the robustness of our findings to potential confounding by unobserved variables, we conducted an e-value analysis (VanderWeele & Ding, 2017). The e-value estimates the minimum strength of association, on the risk ratio scale, that an unobserved confounder would need to have with both the exposure and the outcome in order to completely explain away the estimated association. A large e-value suggests that an unobserved variable would need to pose very strong confounding in order to invalidate the findings.

3. Results

3.1. Characteristics of the sample

The demographic characteristics are presented in Table 1. Out of 242 participants, most (75%) were women. Most of the participants (181 [75%]) were children of the patients, 36 (15%) were grandchildren, while 25 (10%) were spouses or others. Caregivers' mean age was 44.4 years (standard deviation [SD], 14.8), and they provided care for a mean duration of 8.4 years (SD, 2.6). Caregiving burden was high, with a mean ZBI of 56.9 (SD, 12.9). Most of the patients had severe impairment of activities of daily living, as suggested by the mean BADLS of 43.6 (SD, 10.2).

Consumption of fruits, vegetables and milk varied in this sample: fruit, 2.54 days per week (SD, 2.35; median, 2; interquartile range [IQR], 0-3); vegetables, 4.18 days per week (SD, 2.29; median, 4; IQR, 2-7); and milk, 0.65 days per week (SD, 1.58; median, 0; IQR, 0-0). Ninety-three (39%) participants consumed green leafy vegetables, 11 (5%) consumed root vegetables, and 19 (8%) consumed other vegetables. The most commonly reported fruit consumed was avocado (70 [29%]). There was one missing observation for type of fruit consumed and one missing observation for type of vegetable consumed.

3.2. Association between dietary diversity and depression symptom severity

To estimate the association between fruit and vegetable consumption and symptoms of depression, we fitted four multivariable regression models (Table 2). The first regression model included fruit types and their consumption frequency as explanatory variables while adjusting for covariates. Consumption of the different fruit types had statistically significant negative associations with depression symptom severity: avocado, $b = -11.4$ (95% confidence interval [CI], -14.2 to -8.66); citrus, $b = -11.2$ (95% CI, -14.3 to -8.13); and jackfruit, $b = -14.2$ (95% CI, -17.9 to -10.6). Other factors that correlated with depression were caregiver

Table 1
Characteristics of the sample (N = 242).

	N	Mean/Prop.	SD
DASS depression subscale		21.0	11.6
DASS anxiety subscale		20.2	10.9
Caregiver age, yrs		44.4	14.8
Caregiver sex (female)		0.75	0.43
Caregiver education, yrs		5.12	4.55
Caregiving duration, yrs		8.44	2.60
ZBI, total score		56.9	12.9
BADLS, total score		43.6	10.2
Patient age, yrs		86.5	8.85
Patient sex (female)		0.83	0.37
<u>Patient's relationship to caregiver</u>			
Grandparent	36	0.15	
Parent	181	0.75	
Spouse/other	25	0.10	
Fruit consumption freq. (d/wk)		2.54	2.35
<u>Fruit consumption, type</u>			
Avocado	70	0.29	
Citrus	16	0.07	
Jackfruit	15	0.06	
None	140	0.58	
Missing	1	0.004	
Vegetable consumption freq. (d/wk)		4.18	2.29
<u>Vegetable consumption, type</u>			
Green Leafy	93	0.38	
Root	11	0.05	
Other	19	0.08	
None	118	0.49	
Missing	1	.004	
Milk consumption freq. (d/wk)		0.65	1.58

Notes: BADLS, Bristol Activities of Daily Living Scale; DASS, Depression, Anxiety, and Stress Scales; SD, standard deviation; ZBI, Zarit Burden Interview

Table 2

Estimated associations between dietary diversity and depression symptom severity.

Variable	Estimated regression coefficient (b)			
	Model 1	Model 2	Model 3	Model 4
<u>Fruit consumption, type</u>				
Avocado	-11.4***			-0.77
Citrus	-11.2***			-0.74
Jackfruit	-14.2***			-4.68*
None	(base)			(base)
<u>Fruit consumption freq.</u>				
Infrequent (0-2 d/wk)	(base)			(base)
Frequent (3-7 d/wk)	-0.69			-0.06
<u>Vegetable consumption, type</u>				
Green Leafy		-15.6***		-14.1***
Root		-16.8***		-14.0***
Other		-16.4***		-14.8***
None		(base)		(base)
<u>Vegetable consumption freq.</u>				
Infrequent (0-4 d/wk)		(base)		(base)
Frequent (5-7 d/wk)		-1.84		-1.91*
<u>Milk consumption freq.</u>				
None			(base)	(base)
Some (1-7 d/wk)			-7.58***	-2.15*
Caregiver age, yrs	0.09	0.05	0.12*	0.04
Caregiver sex (female)	4.59**	2.57*	6.54***	2.28*
Caregiver education, yrs	0.19	0.05	0.21	0.09
ZBI, total score	0.18***	0.13***	0.25***	0.13***
BADLS, total score	0.11	0.03	0.19**	0.03
Patient age, yrs	-0.08	-0.06	-0.12	-0.06
Patient sex (female)	-0.56	-0.30	-0.64	-0.46
<u>Patient's relationship to caregiver</u>				
Grandparent	(base)	(base)	(base)	(base)
Parent	-2.27	-2.62	-2.61	-2.18
Spouse/other	-2.84	-1.12	-3.35	-0.64
Constant	11.9	24.1***	2.03	24.6***

Notes: BADLS, Bristol Activities of Daily Living Scale; ZBI, Zarit Burden Interview; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

sex (female) and caregiving burden. This model explained 52% of the variation in depression symptom severity.

The second regression model included vegetable types and their consumption frequency as explanatory variables while adjusting for covariates. Consumption of the different vegetable types had statistically significant negative associations with symptoms of depression: green leafy vegetables, $b = -15.6$ (95% CI, -18.1 to -13.0); root vegetables, $b = -16.8$ (95% CI, -20.2 to -13.3); and other vegetables, $b = -16.4$ (95% CI, -20.0 to -12.8). This model explained 66% of the variation in depression symptom severity.

The third regression model included the consumption frequency of milk as an explanatory variable while adjusting for covariates. Frequent milk consumption had a statistically significant negative association with depression symptom severity ($b = -7.58$; 95% CI, -10.2 to -4.93). This model explained 37% of the variation in depression.

In the final regression model for depression symptom severity, we included fruit, vegetable, and milk consumption together as explanatory variables while adjusting for covariates. In this analysis, we found that consumption of jackfruits, green leafy vegetables, root vegetables, other vegetables, frequent vegetable consumption, and frequent milk consumption all had a statistically significant negative association with symptoms of depression. This model explained 66% of the variance in depression symptom severity.

3.3. Association between dietary diversity and anxiety symptom severity

To estimate the association between fruit and vegetable consumption and symptoms of anxiety, we fitted four multivariable regression models (Table 3). The first regression model included fruit types and their consumption frequency as explanatory variables while adjusting for covariates. Consumption of the different fruit types had a statistically

Table 3
Estimated associations between dietary diversity and anxiety symptom severity.

Variable	Estimated regression coefficient (b)			
	Model 1	Model 2	Model 3	Model 4
Fruit consumption, type				
Avocado	-10.7**			-1.47
Citrus	-10.2***			-1.13
Jackfruit	-12.4***			-3.83
None	(base)			(base)
Fruit consumption freq.				
Infrequent (0-2 d/wk)	(base)			(base)
Frequent (3-7 d/wk)	-0.02			0.60
Vegetable consumption, type				
Green Leafy		-13.8***		-12.2***
Root		-15.3***		-12.6***
Other		-14.6***		-12.7***
None		(base)		(base)
Vegetable consumption freq.				
Infrequent (0-4d/wk)		(base)		(base)
Frequent (5-7 d/wk)		-2.03*		-2.07*
Milk consumption freq. (d/wk)				
None			(base)	(base)
Some (1-7 d/wk)			-6.61***	-1.82
Caregiver age, yrs	0.10	0.06	0.12*	0.06
Caregiver sex (female)	4.37***	2.57*	6.04***	2.42*
Caregiver education, yrs	0.14	0.02	0.16	0.05
ZBI, total score	0.18***	0.13***	0.25***	0.13***
BADLS, total score	0.11*	0.05	0.19**	0.05
Patient age, yrs	-0.10	-0.09	-0.14	-0.09
Patient sex (female)	-0.09	0.17	-0.24	0.04
Patient's relationship to caregiver				
Grandparent	(base)	(base)	(base)	(base)
Parent	-2.05	-2.18	-2.18	-2.00
Spouse/other	-4.03	-2.20	-4.21	-2.14
Constant	11.6	22.9***	2.90	22.8***

Notes: BADLS, Bristol Activities of Daily Living Scale; ZBI, Zarit Burden Interview; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

significant negative associations with anxiety symptom severity: avocado, $b = -10.7$; (95% confidence interval [CI], -13.3 to -8.18); citrus, $b = -10.2$ (95% CI, -13.0 to -7.47); and jackfruit, $b = -12.4$; (95% CI, -16.1 to -8.65). Other factors that correlated with anxiety were caregiver sex (female) and caregiving burden. This model explained 51% of the variation in anxiety symptom severity.

The second regression model included vegetable types and their consumption frequency as explanatory variables while adjusting for covariates. Consumption of the different vegetable types and frequency of vegetable consumption had statistically significant negative associations with symptoms of anxiety: green leafy vegetables, $b = -13.8$ (95% CI, -16.2 to -11.4); root vegetables, $b = -15.3$ (95% CI, -18.85 to -11.7); other vegetables, $b = -14.6$ (95% CI, -17.8 to -11.3); and frequent consumption of vegetables, $b = -2.03$; (95% CI, -3.82 to -0.25). This model explained 64% of the variation in anxiety symptom severity.

The third regression model included the consumption frequency of milk as an explanatory variable while adjusting for covariates. Frequent milk consumption had a statistically significant negative association with anxiety symptom severity ($b = -6.61$; 95% CI, -9.08 to -4.14). This model explained 37% of the variation in anxiety symptom severity.

In the final regression model for anxiety symptom severity, we included fruit, vegetable, and milk consumption together as explanatory variables while adjusting for covariates. In this analysis, we found that consumption of green leafy vegetables, root vegetables, other vegetables, frequent vegetable consumption, and frequent milk consumption all had statistically significant negative associations with anxiety symptom severity. This model explained 65% of the variance in anxiety symptom severity.

3.4. Sensitivity analysis

We conducted an e-value analysis to determine the robustness of our

findings to potentially unobserved confounding. In the multivariable regression model for depression symptom severity, the e-value associated with the point estimate for green leafy vegetable consumption was 4.61, and the e-value associated with the confidence interval was 3.44. In the multivariable regression model for anxiety symptom severity, the e-value associated with the point estimate was 4.34, and the e-value associated with the confidence interval was 3.23. These suggests that only very strong confounding would be sufficient to completely explain away the estimated associations between green leafy vegetable consumption and either depression or anxiety symptom severity. An unobserved confounder would need to have a strength of association exceeding 3-4, on the risk ratio scale, with both green leafy vegetable consumption and with depression (or anxiety) symptom severity in order to move our estimated regression coefficients to to have a confidence interval that included 0.

4. Discussion

This study sought to estimate the association between the consumption of different types of fruits and vegetables and symptoms of depression and anxiety in a sample of family caregivers of people living with dementia in southwestern Uganda. We found that the consumption of different categories of fruits and vegetables was negatively correlated with symptoms of both depression and anxiety. These findings are in line with previous studies that found an inverse association between vegetable consumption and symptoms of depression and anxiety (Altun, Brown, Szoeko, & Goodwill, 2019; Baharzadeh et al., 2018; Gibson-Smith et al., 2020; Głabska, Guzek, Groele, & Gutkowska, 2020).

Also similar to our findings on the inverse association between fruit and vegetable consumption and symptoms of depression and anxiety, previous studies have found a dose-response relationship between consumption of fruits and vegetable and improvement in mental well-being and life satisfaction (Blanchflower, Oswald, & Stewart-Brown, 2013; Ocean, Howley, & Ensor, 2019; Peltzer & Pengpid, 2017; White, Horwath, & Conner, 2013). Our results are also consistent with previous studies that have found a correlation between better mental health and specific fruits and vegetables such as citrus fruits, carrots, cucumbers, and green leafy vegetables (Brookie et al., 2018; Conner, Brookie, Carr, Mainvil, & Vissers, 2017; Głabska, Guzek, Groele, & Gutkowska, 2020).

Contrary to other studies that estimated an inverse association between fruit consumption and mental health problems in the general population (Adams & Colner, 2008; Allgöwer, Wardle, & Steptoe, 2001), we did not estimate a statistically significant correlation between fruit types and symptoms of anxiety after adjusting for other groups of vegetables. This discrepancy between our findings and previous studies can potentially be explained by the generally low levels of fruit consumption in this region of southwestern Uganda, which relies heavily on a starch-based diet anchored by the starchy banana *matoke*. In addition, whereas vegetables are readily available in this part of the country, many of these fruits are not locally grown and are typically more expensive than vegetables. Future work, including qualitative research, should attempt to better understand this null finding by conducting a more detailed examination of the consumption of fruits and vegetables in Ugandan communities.

Given the clinical and public health importance of fruit and vegetable consumption for mental well-being in the general population (Boehm et al., 2018; Gehlich et al., 2020; Głabska et al., 2020; Sahraian, Ghanizadeh, & Kazemeini, 2015), our findings are particularly important for bridging the research gap in this area to lay the foundation for clinical and policy work. Future work, perhaps longitudinal studies, may focus on the interactions between consumption of different fruits and vegetables to tease out specific impacts of each category of fruit and vegetables. For caregivers of people with dementia specifically, our findings point to potential dietary interventions that can be used to enhance mental well-being in this population that is at high risk of mental health problems (Ainamani et al., 2020a).

Interpretation of our findings is subject to certain limitations. First, the sample was restricted to caregivers of people with ADRD and might not generalize to general population samples. A second important limitation of our study is the cross-sectional design, which does not allow for causal conclusions. Third, dietary intake was assessed retrospectively and may have been subject to misreporting or recall bias. The latter two limitations belong to a general class of concerns about confounding by unobserved variables: for example, poorer people may be less likely to consume fruits and vegetables, more likely to be food insecure, and more likely to report mental health problems (Lund et al., 2010; Tsai, Bangsberg, et al., 2012). Alternatively, people with memory problems may be more likely to under-report fruit and vegetable consumption and more likely to be depressed. However, our e-value analysis suggests that an unobserved confounder (such as socioeconomic status, food insecurity, or memory impairment) would need to have a very strong association with both fruit and vegetable consumption and with mental health problems in order to move our estimated regression coefficient and its confidence interval to include zero. Despite these limitations, our results suggest that fruits and vegetable consumption is negatively correlated with depressive and anxiety symptoms among the caregivers of people living with dementia.

Abbreviations

BADLS	Bristol Activities of Daily Living Scale
LAMICs	Low- and middle-income countries
ZBI	Zarit Burden Interview

Consent for publication

Not applicable

Availability of data and material

The data sets used and analyzed during the current study are available from the corresponding author on request.

Ethics approval and consent to participate

Ethical approval for the study was obtained from the Mbarara University of Science and Technology. We also received clearance to conduct the study from the Uganda National Council for Science and Technology and the Research Secretariat in the President's office.

Funding

HEA was supported through the Innovative Methods and Metrics for Agriculture and Nutrition Action (IMMANA) programme, led by the London School of Hygiene & Tropical Medicine (LSHTM). IMMANA is confounded with UKAid from the UK government and by the Bill & Melinda Gates Foundation (INV-002962/OPP1211308). Under the grant conditions of the foundation, a Creative Commons Attribution 4.0 Generic License has already been assigned to the author Accepted Manuscripts version that might arise from this submission. ACT acknowledges salary support from the U.S. National Institutes of Health R01MH113494.

Authors' contributions

HEA participated in the conception and design of the study, collected the data, performed data analyses, interpreted the data, and drafted the manuscript. WMB participated in the conception of the study, supervised data collection, and revised the manuscript. GZR, ST, VM, and EMB participated in the conception of the study and revised the manuscript. ACT participated in the conception of the study, supervised data analysis, and provided substantial revision of the manuscript. All authors read and approved the final manuscript.

Declaration of Competing Interest

The authors declare that they have no competing interests.

Acknowledgements

We thank the caregivers who participated in this study; the administrators of Reach One Touch One Ministries and Heal Medical Center, for helping us access caregivers of people with dementia; and our motivated research assistants (Shedrack Atuheire, Rebecca Manoti, Ashaba Patience, and Joseph Namanya), for assisting in data collection and cleaning.

References

- Adams, T., & Colner, W. (2008). The Association of Multiple Risk Factors With Fruit and Vegetable Intake Among a Nationwide Sample of College Students. *Journal of American College Health*, 56, 455–461. <https://doi.org/10.3200/JACH.56.44.455-464>
- Ainamani, Alele, P. E., Rukundo, G. Z., Maling, S., Wakida, E. K., Obua, C., & Tsai, A. C. (2020a). Caregiving burden and mental health problems among family caregivers of people with dementia in rural Uganda. *Global Mental Health*, 7, e13. <https://doi.org/10.1017/gmh.2020.7>
- Ainamani, Alele, P. E., Rukundo, G. Z., Maling, S., Wakida, E. K., Obua, C., & Tsai, A. C. (2020b). Caring for people with dementia in rural Uganda: qualitative study of caregiving burden experienced by informal and formal caregivers. *Journal of Global Health Reports*, 4. <https://doi.org/10.29392/001c.12848>
- Ainamani, Bamwerinde, W. M., Rukundo, G. Z., Tumwesigire, S., Kalibwani, R. M., Bikaitwaho, E. M., & Tsai, A. C. (2021). Participation in gardening activity and its association with improved mental health among family caregivers of people with dementia in rural Uganda. *Preventive Medicine Reports*, 23, Article 101412. <https://doi.org/10.1016/j.pmedr.2021.101412>
- Alfakhri, A. S., Alshudukhi, A. W., Alqahtani, A. A., Alhumaid, A. M., Alhathlol, O. A., Almojali, A. I., ... Alaqeel, M. K. (2018). Depression Among Caregivers of Patients With Dementia. *Inquiry*, 55. <https://doi.org/10.1177/0046958017750432>, 46958017750432-46958017750432.
- Allgöwer, A., Wardle, J., & Steptoe, A. (2001). Depressive symptoms, social support, and personal health behaviors in young men and women. *Health Psychology*, 20(3), 223–227.
- Alqahtani, Alshbriqe, A. A., Awwadh, A. A., Alyami, T. A., Alshomrani, M. S., & Alhazzani, A. (2018). Prevalence and Risk Factors for Depression among Caregivers of Alzheimer's Disease Patients in Saudi Arabia. *Neurology Research International*, 2018, Article 2501835. <https://doi.org/10.1155/2018/2501835>
- Altun, A., Brown, H., Szoek, C., & Goodwill, A. M. (2019). The Mediterranean dietary pattern and depression risk: A systematic review. *Neurology, Psychiatry and Brain Research*, 33, 1–10. <https://doi.org/10.1016/j.npbr.2019.05.007>
- Atteih, S., Mellon, L., Hall, P., Brewer, L., Horgan, F., Williams, D., & Hickey, A. (2015). Implications of stroke for caregiver outcomes: findings from the ASPIRE-S study. *International Journal of Stroke*, 10(6), 918–923. <https://doi.org/10.1111/ijvs.12535>
- Baharzadeh, Siassi, F., Qorbani, M., Koohdani, F., Pak, N., & Sotoudeh, G. (2018). Fruits and vegetables intake and its subgroups are related to depression: a cross-sectional study from a developing country. *Annals of General Psychiatry*, 17, 46. <https://doi.org/10.1186/s12991-018-0216-0>
- Balbara, Y. P. S., Verma, R., Sharma, S., & Mathur, S. (2012). A study of predictors of anxiety and depression among stroke patient-caregivers. *Journal of Mid-Life Health*, 3(1), 31–35. <https://doi.org/10.4103/0976-7800.98815>
- Blanchflower, D. G., Oswald, A. J., & Stewart-Brown, S. (2013). Is Psychological Well-Being Linked to the Consumption of Fruit and Vegetables? *Social Indicators Research*, 114(3), 785–801. Retrieved from <http://www.jstor.org/stable/24720280>.
- Boehm, J. K., Soo, J., Zevon, E. S., Chen, Y., Kim, E. S., & Kubzansky, L. D. (2018). Longitudinal associations between psychological well-being and the consumption of fruits and vegetables. *Health Psychology*, 37(10), 959–967. <https://doi.org/10.1037/hea0000643>
- Brookie, K. L., Best, G. I., & Conner, T. S. (2018). Intake of Raw Fruits and Vegetables Is Associated With Better Mental Health Than Intake of Processed Fruits and Vegetables. *Frontiers in Psychology*, 9, 487. <https://doi.org/10.3389/fpsyg.2018.00487>
- Bucks, R. S., Ashworth, D. L., Wilcock, G. K., & Siegfried, K. (1996). Assessment of activities of daily living in dementia: development of the Bristol Activities of Daily Living Scale. *Age and Ageing*, 25(2), 113–120. <https://doi.org/10.1093/ageing/25.2.113>
- Choda, N., Wakai, K., Naito, M., Imaeda, N., Goto, C., Maruyama, K., ... Hamajima, N. (2020). Associations between diet and mental health using the 12-item General Health Questionnaire: cross-sectional and prospective analyses from the Japan Multi-Institutional Collaborative Cohort Study. *Journal of Nutrition*, 19(1), 2. <https://doi.org/10.1186/s12937-019-0515-6>
- Clyburn, L. D., Stones, M. J., Hadjistavropoulos, T., & Tuokko, H. (2000). Predicting caregiver burden and depression in Alzheimer's disease. *The Journals of Gerontology Series B Psychological Sciences and Social Sciences*, 55(1), S2–13. <https://doi.org/10.1093/geronb/55.1.s2>
- Conner, T. S., Brookie, K. L., Carr, A. C., Mainvil, L. A., & Vissers, M. C. (2017). Let them eat fruit! The effect of fruit and vegetable consumption on psychological well-being

- in young adults: A randomized controlled trial. *PLoS One*, 12(2), Article e0171206. <https://doi.org/10.1371/journal.pone.0171206>
- Crawford, J. R., & Henry, J. D. (2003). The Depression Anxiety Stress Scales (DASS): normative data and latent structure in a large non-clinical sample. *British Journal of Clinical Psychology*, 42(Pt 2), 111–131. <https://doi.org/10.1348/014466503321903544>
- Crichton, G. E., Murphy, K. J., & Bryan, J. (2010). Dairy intake and cognitive health in middle-aged South Australians. *Asia Pacific Journal of Clinical Nutrition*, 19(2), 161–171.
- Dotchin, C. L., Paddick, S. M., Longdon, A. R., Kisoli, A., Gray, W. K., Dewhurst, F., & Walker, R. W. (2014). A comparison of caregiver burden in older persons and persons with Parkinson's disease or dementia in sub-Saharan Africa. *International Psychogeriatrics*, 26(4), 687–692. <https://doi.org/10.1017/S10416161021300255X>
- Gehlich, K. H., Beller, J., Lange-Asschenfeldt, B., Köcher, W., Meinke, M. C., & Lademann, J. (2020). Consumption of fruits and vegetables: improved physical health, mental health, physical functioning and cognitive health in older adults from 11 European countries. *Aging & Mental Health*, 24(4), 634–641. <https://doi.org/10.1080/13607863.2019.1571011>
- Gibson-Smith, D., Bot, M., Brouwer, I. A., Visser, M., Giltay, E. J., & Penninx, B. (2020). Association of food groups with depression and anxiety disorders. *European Journal of Nutrition*, 59(2), 767–778. <https://doi.org/10.1007/s00394-019-01943-4>
- Gilbody, S., Lightfoot, T., & Sheldon, T. (2007). Is low folate a risk factor for depression? A meta-analysis and exploration of heterogeneity. *Journal of Epidemiology and Community Health*, 61(7), 631–637. <https://doi.org/10.1136/jech.2006.050385>
- Głowska, Guzdek, Groele, & Gutkowska. (2020). Fruit and Vegetable Intake and Mental Health in Adults: A Systematic Review. *Nutrients*, 12(1). <https://doi.org/10.3390/nu12010115>
- Gloster, A. T., Rhoades, H. M., Novy, D., Klotsche, J., Senior, A., Kunik, M., ... Stanley, M. A. (2008). Psychometric properties of the Depression Anxiety and Stress Scale-21 in older primary care patients. *Journal of Affective Disorders*, 110(3), 248–259. <https://doi.org/10.1016/j.jad.2008.01.023>
- Guo, Y.-J., & Liu, Y.-J. (2015). Family functioning and depression in primary caregivers of stroke patients in China. *International Journal of Nursing Sciences*, 2(2), 184–189. <https://doi.org/10.1016/j.ijnss.2015.05.002>
- Jefferson, A. L., Byerly, L. K., Vanderhill, S., Lambe, S., Wong, S., Ozonoff, A., & Karlawish, J. H. (2008). Characterization of activities of daily living in individuals with mild cognitive impairment. *American Journal of Geriatric Psychiatry*, 16(5), 375–383. <https://doi.org/10.1097/JGP.0b013e318162f197>
- Kabwama, S. N., Bahendeka, S. K., Wesonga, R., Mutungi, G., & Guwatudde, D. (2019). Low consumption of fruits and vegetables among adults in Uganda: findings from a countrywide cross-sectional survey. *Archives of Public Health*, 77, 4. <https://doi.org/10.1186/s13690-019-0332-6>
- Kakongi, N., Rukundo, G. Z., Gelaye, B., Wakida, E. K., Obua, C., & Okello, E. S. (2020). Exploring Pathways to Hospital Care for Patients with Alzheimer's Disease and Related Dementias in Rural South Western Uganda. *BMC Health Services Research*, 20(1), 498. <https://doi.org/10.1186/s12913-020-05365-5>
- Kamoga, R., Rukundo, G. Z., Wakida, E. K., Nakidde, G., Obua, C., & Buss, S. S. (2019). Dementia assessment and diagnostic practices of healthcare workers in rural southwestern Uganda: a cross-sectional qualitative study. *BMC Health Services Research*, 19(1), 1005. <https://doi.org/10.1186/s12913-019-4850-2>
- Kwon, J., Park, E. C., Kim, W., Choi, D. W., & Jang, S. I. (2019). Depressive symptoms in individuals with family members requiring ADL assistance. *Environmental Health and Preventive Medicine*, 24(1), 49. <https://doi.org/10.1186/s12199-019-0804-x>
- Li, Z., Li, B., Song, X., & Zhang, D. (2017). Dietary zinc and iron intake and risk of depression: A meta-analysis. *Psychiatry Research*, 251, 41–47. <https://doi.org/10.1016/j.psychres.2017.02.006>
- Liu, X., Yan, Y., Li, F., & Zhang, D. (2016). Fruit and vegetable consumption and the risk of depression: A meta-analysis. *Nutrition*, 32(3), 296–302. <https://doi.org/10.1016/j.nut.2015.09.009>
- Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour Research and Therapy*, 33(3), 335–343. [https://doi.org/10.1016/0005-7967\(94\)00075-u](https://doi.org/10.1016/0005-7967(94)00075-u)
- Lund, C., Breen, A., Flisher, A. J., Kakuma, R., Corrigall, J., Joska, J. A., ... Patel, V. (2010). Poverty and common mental disorders in low and middle income countries: A systematic review. *Social Science & Medicine*, 71(3), 517–528. <https://doi.org/10.1016/j.socscimed.2010.04.027>
- McMartin, S. E., Jacka, F. N., & Colman, I. (2013). The association between fruit and vegetable consumption and mental health disorders: evidence from five waves of a national survey of Canadians. *Preventive Medicine*, 56(3-4), 225–230. <https://doi.org/10.1016/j.ypmed.2012.12.016>
- Minichil, W., Getinet, W., Derajew, H., & Seid, S. (2019). Depression and associated factors among primary caregivers of children and adolescents with mental illness in Addis Ababa, Ethiopia. *BMC Psychiatry*, 19(1), 249. <https://doi.org/10.1186/s12888-019-2228-y>
- Mubangizi, V., Maling, S., Obua, C., & Tsai, A. C. (2020). Prevalence and correlates of Alzheimer's disease and related dementias in rural Uganda: cross-sectional, population-based study. *BMC Geriatrics*, 20(1), 48. <https://doi.org/10.1186/s12877-020-1461-z>
- Ocean, N., Howley, P., & Ensor, J. (2019). Lettuce be happy: A longitudinal UK study on the relationship between fruit and vegetable consumption and well-being. *Social Science & Medicine*, 222, 335–345. <https://doi.org/10.1016/j.socscimed.2018.12.017>
- Omrani, V., Haghizadeh, E., & Akoucheian, S. (2018). Depression in Main Caregivers of Dementia Patients: Prevalence and Predictors. *Advanced Biomedical Research*, 7. <https://doi.org/10.4103/2277-9175.225924>, 34-34.
- Peltzer, K., & Pengpid, S. (2017). Dietary consumption and happiness and depression among university students: A cross-national survey. *Journal of Psychology in Africa*, 27(4), 372–377. <https://doi.org/10.1080/14330237.2017.1347761>
- Péneau, S., Galan, P., Jeandel, C., Ferry, M., Andreeva, V., Hercberg, S., & the SU.VI.MAX 2 Research Group. (2011). Fruit and vegetable intake and cognitive function in the SU.VI.MAX 2 prospective study. *American Journal of Clinical Nutrition*, 94(5), 1295–1303. <https://doi.org/10.3945/ajcn.111.014712>. %J The American Journal of Clinical Nutrition.
- Sahraian, A., Ghanizadeh, A., & Kazemini, F. (2015). Vitamin C as an adjuvant for treating major depressive disorder and suicidal behavior, a randomized placebo-controlled clinical trial. *Trials*, 16. <https://doi.org/10.1186/s13063-015-0609-1>, 94-94.
- Sallim, Sayamanathan, A., Cuttilan, A., & Ho, R. (2015). Prevalence of Mental Health Disorders Among Caregivers of Patients With Alzheimer Disease. *Journal of the American Medical Directors Association*, 16(12), 1034–1041.
- Smith, A. P., & Rogers, R. (2014). Positive effects of a healthy snack (fruit) versus an unhealthy snack (chocolate/crisps) on subjective reports of mental and physical health: a preliminary intervention study. *Frontiers in Nutrition*, 1, 10. <https://doi.org/10.3389/fnut.2014.00010>, 10-10.
- Springate, B. A., & Tremont, G. (2014). Dimensions of caregiver burden in dementia: impact of demographic, mood, and care recipient variables. *American Journal of Geriatric Psychiatry*, 22(3), 294–300. <https://doi.org/10.1016/j.jagp.2012.09.006>
- Tsai, Bangsberg, D. R., Frongillo, E. A., Hunt, P. W., Muzaora, C., Martin, J. N., & Weiser, S. D. (2012). Food insecurity, depression and the modifying role of social support among people living with HIV/AIDS in rural Uganda. *Social Science & Medicine* (1982), 74(12), 2012–2019. <https://doi.org/10.1016/j.socscimed.2012.02.033>
- Tsai, Chang, T. L., & Chi, S. H. (2012). Frequent consumption of vegetables predicts lower risk of depression in older Taiwanese - results of a prospective population-based study. *Public Health Nutrition*, 15(6), 1087–1092. <https://doi.org/10.1017/s1368980011002977>
- Van der Does, A. J. (2001). The effects of tryptophan depletion on mood and psychiatric symptoms. *Journal of Affective Disorders*, 64(2-3), 107–119. [https://doi.org/10.1016/s0165-0327\(00\)00209-3](https://doi.org/10.1016/s0165-0327(00)00209-3)
- VanderWeele, T. J., & Ding, P. (2017). Sensitivity analysis in observational research: introducing the E-value. *Annals of Internal Medicine*, 167(4), 268–274. <https://doi.org/10.7326/M16-2607>
- White, B. A., Horwath, C. C., & Conner, T. S. (2013). Many apples a day keep the blues away—daily experiences of negative and positive affect and food consumption in young adults. *British Journal of Health Psychology*, 18(4), 782–798. <https://doi.org/10.1111/bjhp.12021>
- Zarit, S. H., Reever, K. E., & Bach-Peterson, J. (1980). Relatives of the impaired elderly: correlates of feelings of burden. *Gerontologist*, 20(6), 649–655. <https://doi.org/10.1093/geront/20.6.649>